


Vaccination Schedule (standard vaccinations) for babies, children, adolescents and adults¹

English



Vaccination	Age in weeks		months										years						
	6	2	3	4	5-10	11*	12	13-14	15	16-23	2-4	5-6	7-8	9-14	15-16	17	from 18	from 60	
Rotaviruses	U4		G1 ^a	G2	(G3)														
Tetanus ^b	G1		G2		G3 ^a							A1		A2		A'			
Diphtheria ^b	G1		G2		G3 ^a							A1		A2		A'			
Pertussis ^b	G1		G2		G3 ^a							A1		A2		A3'			
Hib ^b – H. influenzae Type b	G1		G2		G3 ^a														
Poliomyelitis ^b	G1		G2		G3 ^a							A1							
Hepatitis B ^b	G1		G2		G3 ^a														
Pneumococcal disease ^b	G1		G2		G3 ^a														S
NEW Meningococcal disease B ^b	G1		G2					G3 ^a											
Meningococcal disease C							G1												
Measles						G1		G2										S ^c	
Mumps, rubella						G1		G2											
Varicella						G1		G2											
HPV – human papillomaviruses												G1 ^a G2 ^a							
Herpes zoster																	G1	G2	
Influenza																	S (annually) ^d		
NEW COVID-19																	G1 ^a , G2 ^a , G3 ^a		S (annually) ^d

Recommended time of vaccination

Catch-up vaccination for basic or primary immunisation for persons not yet vaccinated or for completion of an incomplete vaccination series

Explanatory notes

- A Primary immunisation (up to 3 doses of vaccine G1–G3)
- G Booster vaccination
- S Basic vaccination

1. RKI 2024. Epid Bull 2024;4:1-72.

- a The first vaccination dose should be administered from the age of 6 weeks. Depending on the type of vaccination, 2 or 3 doses are required at intervals of at least 4 weeks.
- b Premature infants: additional dose of vaccine at the age of 3 months, i.e. 4 doses of vaccine in total; infants (including premature babies) are vaccinated with PCV13 or PCV15
- c According to the summary of product characteristics, the vaccination series consists of 3 vaccine doses from the age of 2–23 months, and 2 vaccine doses from the age of 24 months
- d Minimum interval since previous dose: 6 months.
- e Two vaccination doses at intervals of at least 5 months; a 3rd dose is necessary between the 1st and the 2nd dose for catch-up vaccinations from the age of ≥ 15 years or if the interval between vaccinations was < 5 months.
- f Td booster vaccination every 10 years. The next due Td vaccination as a single dose vaccination in the form of Tdap or, if indicated, in the form of a Tdap-IPV combination vaccine.

- g Single dose MMR vaccination for all individuals ≥ 18 years with unclear vaccination status born after 1970, who have not been vaccinated or only received one vaccination as a child.
- h Vaccination until the number of ≥ 3 SARS-CoV-2 antigen contacts required for basic immunity (of which at least 1 vaccination) is reached. Minimum vaccination interval between G1 and G2 ≥ 4 to preferably 12 weeks, and between G2 and G3 ≥ 6 months
- i Vaccination with PCV20
- j Two vaccinations with the adjuvanted herpes zoster vaccine administered at least 2 but no more than 6 months apart.
- k Annual vaccination in autumn
- * Vaccinations can be spread over several vaccination appointments. MMR and V can be administered on the same date or 4 weeks apart.

MAT-DE-2400751-V1-0-02/2024