

# Vaccination schedule for children, adolescents and adults in the Free State of Saxony. (Status 01.01.2024)



Vaccine#	Post-natal	6 Wks	2 Mos	3 Mos	4 Mos	6 Mos	11 Mos	12 Mos (1 year)	14 Mos	23 Mos	4 Yrs	5 Yrs	9 Yrs	10 Yrs	25 Yrs	≥ 50 Yrs	≥ 60 Yrs	every 10 yrs		
<b>Hepatitis B and Hepatitis A (HBV/HAV)</b>		HBV1/HBV 2 (interval > 4 weeks)						HBV 3/4 or HAV 1 ► HAV 2 (interval 6–12 months) or HAV/HBV 1 ► HAV/HBV 2 (interval > 4 weeks) ► HAV/HBV 3 (interval 6–12 months)												
<b>Diphtheria, tetanus, pertussis *</b>			DTPa 1	DTPa 2	DTPa 3			DTPa 4			DTPa 5			Tdap					Tdap	
<b>Haemophilus influenzae type b *</b>			Hib 1	*	Hib 2			Hib 3												
<b>Polio (IPV) * (trivalent)</b>			IPV 1	*	IPV 2			IPV 3						IPV 4					IPV	
<b>Measles, mumps, rubella (MMR)</b>								MMR 1	MMR 2				MMR 3 <sup>†</sup>							
<b>Varicella (VZV)</b>								VZV 1	VZV 2											
<b>Meningococcal B</b>			Meningococcal (group B)																	
<b>Meningococcal ACWY/C</b>			Meningococcal (groups ACWY/C)																	
<b>Influenza **</b>								Influenza annually												
<b>COVID-19 **</b>																			COVID-19	
<b>Pneumococcal ***</b>			PCV 1 ► PCV 2 (interval > 2 months), PCV 3 (interval > 7 months)																	Pneumo-coccal
<b>Rotaviruses</b>			Rotaviruses																	
<b>Human papillomaviruses (HPV) ***</b>													≤ 14 y HPV1 ► HPV 2 ≥ 15 y HPV1 ► HPV2 ► HPV3							
<b>Herpes zoster ***</b>																			RZV 1 ► RZV 2 (interval >6 months)	

# It is essential to observe the summary of product characteristics for the respective vaccines and the following tables with information on application and vaccination intervals! \* All susceptible persons (see tables below) \* For antigen combinations containing a pertussis component, three injections are required in infancy; In full-term infants, the second vaccination planned at 3 months of age can be omitted when using 6- or 5-fold vaccines (known as the 2 + 1 schedule). \*\* Annual seasonal booster with a vaccine adapted to the circulating virus variants (COVID-19) or strain (influenza) \*\*\* The following vaccines are recommended as a preference: pneumococcal (from the age of 18): 20-valent conjugate vaccine (PCV20), HPV: 9-valent vaccine; Herpes zoster: recombinant, adjuvanted vaccine (RZV)